

---

**Product Data Sheet**


---

Product Name: Azimilide  
 Cat. No.: GC14515

**Chemical Properties**

Cas. No. 149908-53-2

Chemical Name 1-[(E)-[5-(4-chlorophenyl)furan-2-yl]methylideneamino]-3-[4-(4-methylpiperazin-1-yl)butyl]imidazolidine-2,4-dione

SMILES CN1CCN(CC1)CCCCN2C(=O)CN(C2=O)N=CC3=CC=C(O3)C4=CC=C(C=C4)Cl

Formula  $C_{23}H_{28}ClN_5O_3$  M.Wt 457.95

Solubility <4.58mg/mL in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure

**Background**

Azimilide(NE-10064) is a class III antiarrhythmic compound, inhibits I(Ks) and I(Kr) in guinea-pig cardiac myocytes and I(Ks) (minK) channels expressed in Xenopus oocytes. IC50 value: Target: in vitro: Azimilide blocked HERG channels at 0.1 and 1 Hz with IC50s of 1.4 microM and 5.2 microM respectively. Azimilide blockade of HERG channels expressed in Xenopus oocytes and I(Kr) in mouse AT-1 cells was decreased under conditions of high [K+]e, whereas block of slowly activating I(Ks) channels was not affected by changes in [K+]e [1]. Azimilide suppressed the following currents (Kd in parenthesis): IKr ( or = 50 microM at +50 and -140 mV, respectively). Azimilide blocked IKr, IKs, and INa in a use-dependent manner. Furthermore, azimilide reduced a slowly inactivating component of Na current that might be important for maintaining the action potential plateau in canine ventricular myocytes [2]. In guinea pig ventricular myocytes, NE-10064 (0.3-3 microM) significantly prolonged action potential duration (APD) at 1 Hz. At 3 Hz, NE-10064 (0.3-1 microM) increased APD only slightly, and at 10 microM decreased APD and the plateau potential. NE-10064 potently blocked the rapidly

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

---

## Product Data Sheet

---

activating component of the delayed rectifier, IKr (IC<sub>50</sub> 0.4 microM), and inhibited IKs (IC<sub>50</sub> 3 microM) with nearly 10-fold less potency [3]. *in vivo*: NE-10064 (10 mg/kg intravenously, i.v.) reduced ( $p < 0.05$ ) the incidence (8 of 12) of PES-induced ventricular tachycardia (VT). The cycle length of induced VT was not prolonged by NE-10064 (0.245 +/- 0.046 s predrug vs. 0.301 +/- 0.060 s postdrug). NE-10064 increased ventricular effective refractory period (VERP 166 +/- 5 ms predrug vs. 194 +/- 13 ms postdrug,  $p = 0.013$ ), prolonged QTc interval (310 +/- 12 ms predrug vs. 350 +/- 16 ms postdrug,  $p = 0.004$ ) and prolonged the effective refractory period (ERP) of noninfarcted myocardium ( $p = 0.045$ ) [4].

### References:

- [1]. Busch AE, et al. Blockade of HERG channels by the class III antiarrhythmic azimilide: mode of action. *Br J Pharmacol.* 1998 Jan;123(1):23-30.
- [2]. Yao JA, et al. Azimilide (NE-10064) can prolong or shorten the action potential duration in canine ventricular myocytes: dependence on blockade of K, Ca, and Na channels. *J Cardiovasc Electrophysiol.* 1997 Feb;8(2):184-98.
- [3]. Fermini B, et al. Use-dependent effects of the class III antiarrhythmic agent NE-10064 (azimilide) on cardiac repolarization: block of delayed rectifier potassium and L-type calcium currents. *J Cardiovasc Pharmacol.* 1995 Aug;26(2):259-71.
- [4]. Black SC, et al. Protection against programmed electrical stimulation-induced ventricular tachycardia and sudden cardiac death by NE-10064, a class III antiarrhythmic drug. *J Cardiovasc Pharmacol.* 1993 Dec;22(6):810-8.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA