
Product Data Sheet

Product Name: DAPTA
Cat. No.: GC17040

Chemical Properties

Cas. No. 106362-34-9

Formula $C_{35}H_{56}N_{10}O_{15}$ M.Wt 856.89

Solubility Soluble to 1 mg/ml in sterile water Storage Desiccate at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure

Protocol

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Kinase experiment:

A novel FITC-labeled tracer from soluble gp120 proteins (25 g/mL) is prepared using a Fluorescent protein labeling kit, according to the manufacture's instructions. Uncoupled FLUOS is removed by Sephadex G-10 column filtration. The molar ratio between FLUOS-labeling molecules and protein is from 3.5 to 4.5 fluorescence molecules per molecule of gp120. The concentration of fluorescent-labeled proteins is measured by Bradford assay and Western blotting by using calibrating amounts of soluble molecules with known concentration. Binding assays are performed in binding buffer, in final volume 100 μ l. Binding is carried out for 1 h at 37°C in 96-well filter plates. Unbound-labeled proteins are removed by rapid vacuum filtration and washing using a 96-well plates manifold. Each binding mix is washed five times with 0.2 mL (total volume of 1.0 mL/well) cold washing buffer (50 mM HEPES, pH 7.4, 150 mM NaCl, 5 mM MgCl₂, 1 mM CaCl₂). Filters are counted with a fluorescent plate reader at 495/530 nm.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1]. Pollicita M, et al. Profound anti-HIV-1 activity of DAPTA in monocytes/macrophages and inhibition of CCR5-mediated apoptosis in neuronal cells. *Antivir Chem Chemother.* 2007;18(5):285-95.
- [2]. Polianova MT, et al. Chemokine receptor-5 (CCR5) is a receptor for the HIV entry inhibitor peptide T (DAPTA). *Antiviral Res.* 2005 Aug;67(2):83-92.

Background

DAPTA is a synthetic peptide, functions as a viral entry inhibitor by targeting selectively CCR5, and shows potent anti-HIV activities.

DAPTA (1 nM) inhibits HIV-1 replication in monocytes/macrophages (M/M) by >90%. DAPTA blocks HIV entry and prevents HIV-1 infection. DAPTA reduces CCR5 mAb binding in human primary macrophages. DAPTA potently blocks R5 gp120-mediated neuronal apoptosis. DAPTA is even more potent in preventing neuronal apoptosis than the CCR5 antagonist TAK-779[1]. DAPTA potently inhibits specific CD4-dependent binding of gp120 BaL (IC₅₀ = 0.06 nM) and CM235 (IC₅₀ = 0.32 nM) to CCR5. DAPTA (1 nM) blocks formation of the gp120/sCD4 complex with CCR5. DAPTA inhibits the binding of gp120BaL/sCD4 to CCR5 (Cf2Th/synR5) cells with IC₅₀ of 55 ± 0.08 pM[2].

References:

- [1]. Pollicita M, et al. Profound anti-HIV-1 activity of DAPTA in monocytes/macrophages

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

and inhibition of CCR5-mediated apoptosis in neuronal cells. Antivir Chem Chemother. 2007;18(5):285-95.

[2]. Polianova MT, et al. Chemokine receptor-5 (CCR5) is a receptor for the HIV entry inhibitor peptide T (DAPTA). Antiviral Res. 2005 Aug;67(2):83-92.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA