

---

**Product Data Sheet**

---

Product Name: Cadazolid

Cat. No.: GC19085

**Chemical Properties**

Cas. No. 1025097-10-2

SMILES O=C(C1=CN(C2CC2)C3=C(C=C(F)C(N4CCC(O)(COC5=CC=C(N6C(O[C@@H](CO)C6)=O)C=C5F)CC4)=C3)C1=O)OFormula C<sub>29</sub>H<sub>29</sub>F<sub>2</sub>N<sub>3</sub>O<sub>8</sub>

M.Wt 585.55

Solubility DMSO : ≥ 150 mg/mL (256.17 mM)

Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure **Protocol****Cell experiment:**

MICs of cadazolid, metronidazole, vancomycin, moxifloxacin and linezolid are determined using agar incorporation for 100 Clostridium difficile isolates, including 30 epidemic strains (ribotypes 027, 106 and 001) with reduced metronidazole susceptibility, 2 linezolid-resistant isolates and 2 moxifloxacin-resistant isolates. The efficacy of two cadazolid dosing regimens (250 versus 750 mg/L twice daily for 7 days) to treat simulated CDI is evaluated. Microflora populations, Clostridium difficile total viable counts and spores, cytotoxin titres, possible emergence of cadazolid, linezolid or quinolone resistance, and antimicrobial concentrations are monitored throughout[2].

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

### References:

- [1]. Baldoni D, et al. Cadazolid, a novel antibiotic with potent activity against *Clostridium difficile*: safety, tolerability and pharmacokinetics in healthy subjects following single and multiple oral doses. *J Antimicrob Chemother.* 2014 Mar;69(3):706-14.
- [2]. Chilton CH, et al. In vitro activity of cadazolid against clinically relevant *Clostridium difficile* isolates and in an in vitro gut model of *C. difficile* infection. *J Antimicrob Chemother.* 2014 Mar;69(3):697-705.

### Background

Cadazolid (ACT-179811) is a new oxazolidinone antibiotic with potent activity against *Clostridium difficile*.

Cadazolid is a new antibiotic in development for the treatment of *Clostridium difficile*-associated diarrhea[1]. Cadazolid is active against all (including linezolid- and moxifloxacin-resistant) *Clostridium difficile* strains (MIC90 0.125, range 0.03-0.25 mg/L).

**Caution: Product has not been fully validated for medical applications. For research use only.**

**Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com**

**Address: 10292 Central Ave. #205, Montclair, CA, USA**

---

## Product Data Sheet

---

The cadazolid geometric mean MIC is 152-fold, 16-fold, 9-fold and 7-fold lower than those of moxifloxacin, linezolid, metronidazole and vancomycin, respectively. Both cadazolid dosing regimens rapidly reduce *Clostridium difficile* viable counts and cytotoxin with no evidence of recurrence. Cadazolid levels persists at 50-100-fold supra-MIC for 14 days post-dosing. Cadazolid inhibition of enumerated gut microflora is limited, with the exception of bifidobacteria; *Bacteroides fragilis* group and *Lactobacillus* spp. counts are unaffected. There is no evidence for selection of strains resistant to cadazolid, quinolones or linezolid[2].

Cadazolid is well tolerated up to 3000 mg given twice daily for 10 days. The most common adverse event is headache, with no observed relationship between dose or treatment duration and adverse events. Plasma concentrations of cadazolid are low. No plasma concentrations >3.3 ng/mL are observed after single doses or >6.9 ng/mL after 10 days of multiple doses. Food increased the mean C<sub>max</sub> from 0.73 to 1.87 ng/mL and mean AUC<sub>0-t</sub> from 3.13 to 15.69 ng.h/mL after a single 300 mg dose. The increase in systemic exposure to cadazolid across doses is less than dose-proportional. The mean cumulative faecal recovery is 81.0%-93.5%. Urinary recovery of unchanged compound is less than 0.015%[1].

### References:

- [1]. Baldoni D, et al. Cadazolid, a novel antibiotic with potent activity against *Clostridium difficile*: safety, tolerability and pharmacokinetics in healthy subjects following single and multiple oral doses. *J Antimicrob Chemother.* 2014 Mar;69(3):706-14.
- [2]. Chilton CH, et al. In vitro activity of cadazolid against clinically relevant *Clostridium difficile* isolates and in an in vitro gut model of *C. difficile* infection. *J Antimicrob Chemother.* 2014 Mar;69(3):697-705.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA