
Product Data Sheet

Product Name: S-8510 phosphate (SB-737552 phosphate)

Cat. No.: GC31121

Chemical Properties

Cas. No. 151466-23-8

SMILES O=P(O)(O)O.C12=NC=C3C(NC(C4=NOC=C4)=N3)=C1COCC2Formula $C_{12}H_{13}N_4O_6P$ M.Wt 340.23

Solubility Soluble in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure **Protocol**

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Mice, Rats[1] Proconvulsant actions of S-8510 are examined in combination with a subconvulsive dose of convulsant (PTZ or strychnine) or minimal electroconvulsive shock in ddY mice. Each convulsant is administered, simultaneously with the inverse agonist, whereas minimal electroconvulsive shock is applied 5 min after intravenous injection of the inverse agonist. The proconvulsant test of S-8510 on PTZ (52, 62.5, 75 or 90 mg/kg s.c.) or strychnine (0.25 mg/kg, s.c.)-induced convulsion is made. ED50 values for producing clonic convulsion to death in 50% of ddY mice are calculated. Anxiogenic actions of each inverse agonist are examined in the punished water licking method in Wistar rats. Each inverse agonist is administered per OS 30 min before the test trial. Antiamnesic actions of inverse agonists are examined in the water maze paradigm. Wistar rats are given 4 trials every day for consecutive 4 days. On the 5th day, each animal has the retention test for 60 s. Amnesia is produced by intraperitoneally injected scopolamine at a dose of 0.5 mg/kg. Scopolamine and each inverse agonist are given 15 min and 30 min before the retention trial, respectively[1].

Animal experiment:

References:

[1]. Kawasaki K, et al. A novel benzodiazepine inverse agonist, S-8510, as a cognitive enhancer. *Prog Neuropsychopharmacol Biol Psychiatry*. 1996 Nov;20(8):1413-25.

Background

S-8510 (phosphate) is an inverse Benzodiazepine (BDZ) receptor agonist, with K_{is} of

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

34.6 nM, 36.2 nM for -GABA and +GABA respectively.

S-8510 has a relatively high affinity to BDZ receptors. The ratio of K_i values for each ligand with and without GABA is defined as the GABA ratio, which is considered as a biochemical index for BDZ receptor ligands. The GABA ratio for S-8510 or CGS8216 is close to the value for flumazenil which is considered as an antagonist or a very weak agonist. S-8510 (10^{-7} M) enhances LTP and this enhancement is antagonized by BDZ receptor antagonist, flumazenil. Flumazenil itself does not affect LTP or evokes responses prior to tetanic stimulation. S-8510 has no effect on the field evoked potential up to 10^{-5} M. However, S-8510 increases the amplitude of the population spike at a dose of 10^{-4} M, and this effect is completely antagonized by concomitant application of flumazenil (10^{-4} M) [1].

S-8510 or CGS8216 could cause lethal convulsion only in combination with more than 90 mg/kg of PTZ. The proconvulsant activity of S-8510 appears to be selective for PTZ-induced subconvulsive state. Scopolamine decreases the time spending in the area around the platform, indicating the amnesic action of scopolamine. S-8510 and CGS8216 reverses this scopolamine-induced amnesia. S-8510 improves the memory impairment induced by diazepam in the water maze and passive avoidance paradigms as well. S-8510 dose-dependently increases the ACh level up to 100 mg/kg. Both S-8510 and PTZ increases the extracellular level of NA in the hippocampus in a dose-dependent manner. Anxiogenic actions of S-8510, CGS8216 and FG7142 are examined in the water lick conflict paradigm of Wistar rats. S-8510 and CGS8216 fail to affect this behavioral paradigm up to 30 mg/kg. S-8510 significantly decreases the immobility time in the forced swimming test using ddY mice at 40 to 80 mg/kg in a dose-dependent manner. In the tetrabenazine-induced ptosis model, S-8510 significantly reduces the extent of ptosis induced by tetrabenazine at doses more than 10 mg/kg. Again, S-8510 reduces the extent of ptosis only by 39% even at the maximum dose, whereas imipramine exerts more pronounced effects (by about 80% at a dose of 20 mg/kg)[1].

[1]. Kawasaki K, et al. A novel benzodiazepine inverse agonist, S-8510, as a cognitive enhancer. *Prog Neuropsychopharmacol Biol Psychiatry*. 1996 Nov;20(8):1413-25.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA