

---

**Product Data Sheet**

---

Product Name: PSI-352938 (PSI-938)

Cat. No.: GC33971

**Chemical Properties**

Cas. No. 1231747-17-3

SMILES O=P1(OC(C)C)OC[C@]2([H])[C@@]([C@@](C)(F)[C@H](N3C(N=C(N)N=C4OCC)=C4N=C3)O2)([H])O1Formula C<sub>16</sub>H<sub>23</sub>FN<sub>5</sub>O<sub>6</sub>P M.Wt 431.36

Solubility Soluble in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure **Protocol**

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

### Cell experiment:

GT 1a, 1b, and 2a replicon cells are cultured in the presence of G418 (0.75 mg/mL for GT 1a, 0.25 mg/mL for GT 1b and 2a) and increasing concentrations of PSI-352938 (PSI-938) or PSI-353661 starting at their respective EC50 or EC90. As a no-compound control, replicon cells are maintained in parallel in the equivalent percent volume (0.2%) of DMSO. Cells are passaged whenever they reach ~80% confluence and replenished with G418 medium containing fresh compound. At various passages, cells are tested for sensitivity to PSI-352938 (PSI-938) and PSI-353661. For each assay, 3-fold dilutions of test compound are added to cells in duplicate and incubated at 37°C in a humidified 5% CO<sub>2</sub> atmosphere for 4 days. Inhibition of HCV replicon RNA replication is determined by real-time PCR (RT-PCR) using primers that anneal to the 5' untranslated region or by measuring the levels of luminescence expressed via the firefly or Renilla luciferase reporter gene using the Bright-Glo or Renilla-Glo reagent, respectively. EC50 and EC90, the concentrations at which 50% and 90% inhibition are achieved, are determined using GraphPad Prism software. Aliquots of cells are also saved for RNA isolation, cDNA synthesis, and PCR amplification for sequencing analysis[1].

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

### References:

- [1]. Lam AM, et al.  
Hepatitis C virus  
nucleotide  
inhibitors PSI-  
352938 and PSI-  
353661 exhibit a  
novel mechanism  
of resistance  
requiring multiple  
mutations within  
replicon RNA. J  
Virol. 2011  
Dec;85(23):12334-  
42.
- [2]. Lam AM, et al.  
Inhibition of  
hepatitis C virus  
replicon RNA  
synthesis by PSI-  
352938, a cyclic  
phosphate  
prodrug of  $\beta$ -D-2'-  
deoxy-2'- $\alpha$ -fluoro-  
2'- $\beta$ -C-  
methylguanosine.  
Antimicrob Agents  
Chemother. 2011  
Jun;55(6):2566-75.

### Background

PSI-352938 (PSI-938) is a hepatitis C virus (HCV) nucleotide inhibitor.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

PSI-352938 (PSI-938) and PSI-353661 inhibit HCV genotype (GT) 1b replicon replication with 50% effective concentrations (EC50s) of  $0.13 \pm 0.076 \mu\text{M}$  and  $3.0 \pm 1.4 \text{ nM}$ , respectively, and are similarly active against GT 1a and 2a replicons and infectious viruses. Metabolism of PSI-352938 and PSI-353661 generates the same 5'-triphosphate metabolite, PSI-352666, which is similarly active against NS5B polymerases from GT 1 to 4[1]. PSI-352938 (PSI-938) is a novel cyclic phosphate prodrug of  $\beta$ -D-2'-deoxy-2'- $\alpha$ -fluoro-2'- $\beta$ -C-methylguanosine 5'-monophosphate that has potent activity against HCV. PSI-352938 (PSI-938) has similar activity against genotype 1a, 1b, and 2a replicons, with EC50s ranging from 0.13 to 0.20  $\mu\text{M}$  and EC90 values ranging from 0.35 to 0.74  $\mu\text{M}$ . PSI-352938 (PSI-938) also effectively inhibits HCV replication in the infectious virus assays: the EC50 and EC90 values are  $0.28 \pm 0.083 \mu\text{M}$  and  $0.63 \pm 0.018 \mu\text{M}$ , respectively, against the H77 infectious virus and  $0.39 \pm 0.31 \mu\text{M}$  and  $1.16 \pm 0.64 \mu\text{M}$ , respectively, against the JFH-1 infectious virus. In contrast, PSI-352938 is not active against HBV or HIV up to the highest concentration tested (EC50 > 100  $\mu\text{M}$ )[2].

[1]. Lam AM, et al. Hepatitis C virus nucleotide inhibitors PSI-352938 and PSI-353661 exhibit a novel mechanism of resistance requiring multiple mutations within replicon RNA. *J Virol.* 2011 Dec;85(23):12334-42. [2]. Lam AM, et al. Inhibition of hepatitis C virus replicon RNA synthesis by PSI-352938, a cyclic phosphate prodrug of  $\beta$ -D-2'-deoxy-2'- $\alpha$ -fluoro-2'- $\beta$ -C-methylguanosine. *Antimicrob Agents Chemother.* 2011 Jun;55(6):2566-75.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA