

---

## Product Data Sheet

---

Product Name: Tubulysin C

Cat. No.: GC37837

**Chemical Properties**

Cas. No. 205304-88-7

SMILES O=C([C@@]([C@@H](C)CC)([H])NC([C@@H](CCCC1)N1C)=O)N(COC(CC)=O)[C@@H](C(C)C)C[C@H](C2=NC(C(N[C@H](C[C@H](C)C(O)=O)CC3=CC=C(O)C=C3)=O)=CS2)OC(C)=O

Formula C<sub>41</sub>H<sub>61</sub>N<sub>5</sub>O<sub>10</sub>S M.Wt 816.02

Solubility Soluble in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure **Background**

Tubulysin C is a highly cytotoxic peptide isolated from the myxobacterial species *Archangium geophyra* and *Angiococcus disciformis*. Tubulysin displays extremely potent cytotoxic activity in mammalian cells, including multidrug-resistant cell lines, with IC<sub>50</sub> values in the lower nanomolar range[1]. Tubulysin C is a cytotoxic activity tubulysin which inhibits tubulin polymerization and leads to cell cycle arrest and apoptosis[2].

[1]. Kubicek K, et al. The tubulin-bound structure of the antimetabolic drug tubulysin. *Angew Chem Int Ed Engl.* 2010 Jun 28;49(28):4809-12. [2]. Vlahov IR, et al. Acid mediated formation of an N-acyliminium ion from tubulysins: a new methodology for the synthesis of natural tubulysins and their analogs. *Bioorg Med Chem Lett.* 2011 Nov 15;21(22):6778-81.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---