

---

**Product Data Sheet**

---

Product Name: Xenin  
Cat. No.: GC37943

**Chemical Properties**

Cas. No. 144092-28-4

Formula  $C_{139}H_{224}N_{38}O_{32}S$  M.Wt 2971.57

Solubility Soluble in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure

**Protocol****Animal experiment:**

Mice: Mice (wild-type and ob/ob mice) are fasted overnight (1800–1000 h) and injected with xenin intracerebroventricularly (0.1, 1, or 5 µg) or intraperitoneally (0.5, 5, 15, or 50 µg/g body wt) at 1000 h. To compare the feeding-suppressing effect between xenin and neurotensin, equimolar amounts (16.5 nmol) of xenin (50 µg/g body wt) or neurotensin (28 µg/g body) are injected intraperitoneally after an overnight fast. Control mice receive either intracerebroventricular injection of aCSF or intraperitoneal injection of saline. Preweighed food is provided to mice immediately after the injection. Cumulative food intake is measured at time points indicated in each figure up to 24 h after injection[2].

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

### References:

- [1]. Cooke JH, et al. Peripheral and central administration of xenin and neurotensin suppress food intake in rodents. Obesity (Silver Spring). 2009 Jun;17(6):1135-43.
- [2]. Leckstrom A, et al. Xenin, a gastrointestinal peptide, regulates feeding independent of the melanocortin signaling pathway. Diabetes. 2009 Jan;58(1):87-94.
- [3]. Kim ER, et al. Xenin delays gastric emptying rate and activates the brainstem in mice. Neurosci Lett. 2010 Aug 30;481(1):59-63.

### Background

Xenin is a 25-amino acid peptide initially isolated from human gastric mucosa. Xenin is a gut hormone that can reduce food intake.

Xenin is abundantly expressed in gastric, duodenal, and jejunal mucosa, and is found at lower levels in the pancreas. Xenin is released into the circulation postprandially and has

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

been reported to stimulate pancreatic endocrine and exocrine secretion, inhibit gastrin secretion, and influence gastrointestinal motility. Xenin is highly homologous to neurotensin. Xenin and neurotensin are reported to have similar biological effects[1].

Both intracerebroventricular and intraperitoneal administration of xenin inhibit fasting-induced hyperphagia in wild-type mice in a dose-dependent manner. The intraperitoneal injection of xenin also reduces nocturnal intake in ad libitum-fed wild-type mice. The intraperitoneal injection of xenin increases Fos immunoreactivity in hypothalamic nuclei, including the paraventricular nucleus and the arcuate nucleus. Xenin reduces food intake in agouti and ob/ob mice[2]. Gastric emptying rate is reduced by about 93% in xenin-treated mice compared to saline-treated control mice. The i.p. xenin injection significantly increases Fos-immunoreactive cells in the nucleus of the solitary tract of the brainstem, but not area postrema and dorsal motor nucleus of the vagus[3].

### References:

- [1]. Cooke JH, et al. Peripheral and central administration of xenin and neurotensin suppress food intake in rodents. *Obesity (Silver Spring)*. 2009 Jun;17(6):1135-43.
- [2]. Leckstrom A, et al. Xenin, a gastrointestinal peptide, regulates feeding independent of the melanocortin signaling pathway. *Diabetes*. 2009 Jan;58(1):87-94.
- [3]. Kim ER, et al. Xenin delays gastric emptying rate and activates the brainstem in mice. *Neurosci Lett*. 2010 Aug 30;481(1):59-63.

**Caution: Product has not been fully validated for medical applications. For research use only.**

**Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com**

**Address: 10292 Central Ave. #205, Montclair, CA, USA**