
Product Data Sheet

Product Name: (S)-(+)-Linoleyl-2'-Hydroxy-1'-Propylamide

Cat. No.: GC41731

Chemical Properties

Cas. No.

SMILES CCCCC/C=C\C/C=C\CCCCCCCC(NC[C@H](C)O)=OFormula $C_{21}H_{39}NO_2$

M.Wt 337.5

Solubility	DMF: 11 mg/ml, DMF: 11 mg/ml, DMSO: 5 mg/ml, DMSO: 5 mg/ml, Ethanol: 20 mg/ml, Ethanol: 20 mg/ml, Ethanol: PBS (pH 7.2)(1:2): 0.3 mg/ml, Ethanol: PBS (pH 7.2)(1:2): 0.3 mg/ml	Store Storage at - 20°C
------------	--	-------------------------------

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure

Background

N-Acyl ethanolamines (NAEs) have diverse biological actions that are strongly affected by the associated acyl group. Linoleoyl ethanolamide (LOEA) has potential signaling roles in aging and neurological functioning. LOEA has a weak affinity for cannabinoid (CB) receptors ($K_i = 10, 25 \mu\text{M}$ for CB1, CB2, respectively) and inhibits voltage-gated K^+ channels. LOEA also inhibits fatty acid amide hydrolase (FAAH; $K_i = 9 \mu\text{M}$) and is hydrolyzed by FAAH. (S)-(+)-Linoleyl-2'-hydroxy-1'-propylamide is a homolog of LOEA, characterized by the addition of an (S)- β -methyl group at the terminal ethanolamine carbon. A similar modification of arachidonoyl ethanolamide to produce S-2-methanandamide imparts higher affinity for the CB receptor. The physiological actions of this compound have not been evaluated.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA
