
Product Data Sheet

Product Name: Elsubrutinib

Cat. No.: GC64819

Chemical Properties

Cas. No. 1643570-24-4

Formula C₁₇H₁₉N₃O₂ M.Wt 297.35

Solubility Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure

Background

Elsubrutinib (ABBV-105) is an orally active, potent, selective and irreversible Bruton's tyrosine kinase (BTK) inhibitor. The IC₅₀ of Elsubrutinib for BTK catalytic domain is 0.18 μM. Elsubrutinib can be used for the research of inflammatory disease[1].

Elsubrutinib inhibits BTK (C481S) with an IC₅₀ of 2.6 μM, indicating a significant loss in potency upon exchanging the targeted thiol nucleophile with an alcohol, suggesting Cys481 is important in the manner in which Elsubrutinib inhibits BTK. Elsubrutinib irreversibly inhibits BTK enzyme activity and blocks BTK-dependent cellular activation. Elsubrutinib inhibits histamine release from IgE-stimulated basophils and IL-6 release from IgG-stimulated monocytes, which utilize Fcε and Fcγ receptors respectively. Elsubrutinib inhibits IgM-mediated B cell proliferation, which is dependent on signaling through the BCR. Elsubrutinib also inhibits TNF-release from CpG-DNA stimulated PBMCs, which signals through TLR9, although it does not inhibit the function of TLRs that do not use ITAM motifs, namely, TNF release from PBMCs stimulated either through TLR4 (with LPS) or through TLR7/8 (with R848). Elsubrutinib has significant impacts on IgM-mediated B cell proliferation[1].

Elsubrutinib (10 mg/kg; p.o.) inhibits antibody responses to NP-Ficoll and NP-KLH, but not to NP-LPS or Prevnar-13[1]. Elsubrutinib (0.1~10 mg/kg; p.o.) results in dose-dependent

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

inhibition of paw swelling throughout the course of disease and significantly prevents the onset of proteinuria and prolongs survival at the 10 mg/kg QD and BID doses, while lower doses does not significantly inhibit these endpoints[1].Elsubrutinib demonstrates exposure-dependent inhibition of increases in paw volume. Elsubrutinib significantly inhibits bone volume loss in a dose dependent manner consistent with the observed anti-inflammatory effects[1].

[1]. Goess C, et al. ABBV-105, a selective and irreversible inhibitor of Bruton's tyrosine kinase, is efficacious in multiple preclinical models of inflammation [published correction appears in Mod Rheumatol. 2019 May;29(3):v]. Mod Rheumatol. 2019;29(3):510-522.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA