
Product Data Sheet

Product Name: FGF 23 C-term Human
 Cat. No.: GP20255
 Batch No.: 1

Product Data

Purity	>98%	Source	E. coli
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	Tumor-derived hypophosphatemia-inducing factor; HYPF; ADHR; HPDR2; PHPTC; FGF23; FGF-23; Fibroblast Growth Factor-23.		
Amino Acid Sequence	MKHHHHHHAS AEDDSERDPL NVLKPRARMT PAPASCSQEL PSAEDNSPMA SDPLGVVRGG RVNTHAGGTG PEGCRPFAKF I.		
Solubility	It is recommended to add deionized water to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it on cell culture.		
Formulation	FGF-23 C-term was filtered (0.4µm) and lyophilized from 0.5mg/ml supplied in 20mM TRIS and 50mM NaCl, pH 7.5.		

Introduction

FGF-23 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-23 inhibits renal tubular phosphate transport. This gene was identified by its mutations associated with autosomal dominant hypophosphatemic rickets (ADHR), an inherited phosphate wasting disorder. Abnormally high level expression of FGF23 was found in oncogenic hypophosphatemic osteomalacia (OHO), a phenotypically similar disease caused by abnormal phosphate metabolism. Mutations FGF23 have also been shown to cause familial tumoral calcinosis with hyperphosphatemia.

Stability

Store lyophilized FGF 23 C-term at -20°C . Aliquot the product after reconstitution to

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

avoid repeated freezing/thawing cycles. Reconstituted FGF 23 C-term can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C .

Background

FGF-23 C-term Protein is 8.67 kDa protein containing 72 amino acid residues and an additional 9 a.a. His-Tag at N-terminus.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA