
Product Data Sheet

Product Name: IL-4 Porcine
 Cat. No.: GP20505
 Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	BCGF; BCDF; B cell stimulating factor; BSF-1; Lymphocyte stimulatory factor 1; IL-4; MGC79402; Binetrakin; Pitrakinra.		
Amino Acid Sequence	MHKCDITLQE IIKTLNILTA RKNSCMELPV TDVFAAPENT TEKETFCRAS TVLRHIYRHH TCMKSLLSGL DRNLSSMANM TCSVHEAKKS TLKDFLERLK TIMKEKYSKC.		
Solubility	It is recommended to reconstitute the lyophilized IL-4 in sterile 18MΩ-cm H ₂ O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.		
Formulation	The protein was lyophilized from a sterile (0.2μm) filtered solution containing 10 mM sodium phosphate, pH 7.5.		

Introduction

IL4 is a pleiotropic cytokine produced by activated T cells. IL4 is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which contributes to various overlapping roles of this cytokine and IL13. STAT6, a signal transducer and activator of transcription plays a main role in mediating the immune regulatory signal of this cytokine. IL4, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. IL4, IL13 and IL5 are regulated co-ordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome.

Biological Activity

The ED₅₀, as determined by TF-1 cell proliferation is <2ng/ml corresponding to a specific activity which is $\geq 5.0 \times 10^5$ units/mg.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Stability

Lyophilized IL-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C . Upon reconstitution IL-4 should be stored at 4°C between 2-7 days and for future use below -18°C . Please prevent freeze-thaw cycles.

Background

Interleukin-4 Porcine Recombinant produced in E. coli is a non-glycosylated monomer chain containing 110 amino acids and having a molecular mass of 12.7kDa. IL-4 is purified by proprietary chromatographic techniques.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA