

---

## Product Data Sheet

---

Product Name: IL 16 Human, (121 a.a.)

Cat. No.: GP20576

Batch No.: 1

**Product Data**

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	IL16; Interleukin-16; LCF; Lymphocyte Chemoattractant Factor; prIL-16; IL-16; FLJ16806; FLJ42735; FLJ44234; HsT19289.		
Amino Acid Sequence	SAASASAASD VSVSTAEAT VCTVTLEKMS AGLGFSLEGG KGSLHGDKPL TINRIFKGAA SEQSETVQPG DEILQLGGTA MQGLTRFEAW NIIKALPDGP VTIVIRRKSL QSKETTAAGD S		
Solubility	It is recommended to reconstitute the lyophilized Interleukin-16 in sterile 18MΩ-cm H <sub>2</sub> O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.		
Formulation	IL-16 was lyophilized from a 0.2μm filtered concentrated solution in PBS, pH 7.4.		

**Introduction**

IL-16 is a pleiotropic cytokine that functions as a chemoattractant, a modulator of T cell activation, and an inhibitor of HIV replication. The signaling process of IL-16 is mediated by CD4. The product of this gene undergoes proteolytic processing, which is found to yield two functional proteins. IL-16 functions exclusively attributed to the secreted C-terminal peptide, while the N-terminal product may play a role in cell cycle control. Caspase 3 is reported to be involved in the proteolytic processing of this protein. Two transcript variants encoding different isoforms have been found for this gene. IL-16 stimulates a migratory response in cd4+ lymphocytes, monocytes, and eosinophils. Also induces t-lymphocyte expression of interleukin 2 receptor, ligand for cd4.

**Biological Activity**

Fully biologically active when compared to standard. Determined by its ability to

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

---

## Product Data Sheet

---

chemoattract human CD4+ T-Lymphocytes using a concentration range of 50.0-100.0 ng/ml.

### Stability

Lyophilized Interleukin-16 although stable at room temperature for 3 weeks, should be stored desiccated below  $-18^{\circ}\text{C}$ . Upon reconstitution IL16 should be stored at  $4^{\circ}\text{C}$  between 2-7 days and for future use below  $-18^{\circ}\text{C}$ . For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

### Background

Interleukin-16 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 121 amino acids and having a molecular mass of 12.4 kDa. The IL-16 is purified by proprietary chromatographic techniques.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA