
Product Data Sheet

Product Name: Cyclophilin F Rat
 Cat. No.: GP21530
 Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped with Ice Packs.
Synonyms	Peptidyl-prolyl cis-trans isomerase F; mitochondrial; PPIase F; Cyclophilin D; CyP-D; CypD; Cyclophilin F; Rotamase F; Ppif; Peptidyl-prolyl cis-trans isomerase F; mitochondrial; PPIase.		
Amino Acid Sequence	MGSSHHHHHH SGLVPRGSH MGSCSDGGAR GANSSSQNPL VYLDVGADGQ PLGRVVLELK ADVVPKTAEN FRALCTGEKG FGYKGSTFHR VIPAFMCQAG DFTNHNGTGG KSIYGSRFPD ENFTLKHVGP GVLSMANAGP NTNGSQFFIC TIKTDWLDGK HVVFGHVKEG MDVVKKIESF GSKSGKTSKK IVITDCGQLS.		
Formulation	Cyclophilin F protein solution (1mg/ml) containing Phosphate buffered saline (pH7.4), 10% glycerol and 1mM DTT.		

Introduction

Peptidyl-prolyl cis-trans isomerase F, mitochondrial (Cyclophilin-F) accelerates the folding of proteins. Cyclophilin-F catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and involved in regulation of the mitochondrial permeability transition pore (mPTP). Cyclophilin-F, in cooperation with mitochondrial TP53, is involved in activating oxidative stress-induced necrosis. Cyclophilin-F is also involved in modulation of mitochondrial membrane F1F0 ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels. Furthermore, Cyclophilin-F has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Background

Cyclophilin F Rat Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 200 amino acids (30-206a.a.) and having a molecular mass of 21.2kDa. Cyclophilin F is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA