
Product Data Sheet

Product Name: PLA2G5 Human
 Cat. No.: GP22100
 Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	Calcium-dependent phospholipase A2; EC 3.1.1.4; Phosphatidylcholine 2-acylhydrolase; PLA2-10; Group V phospholipase A2; GV-PLA2; MGC46205; hVPLA(2); DKFZp686C2294; sPLA2-V; PLA2G5.		
Amino Acid Sequence	MRGSHHHHHH GMASHMGLLD LKSMIEKVTG KNALTNYGFY GCYCGWGGRG TPKDGTDWCC WAHDHCYGRLEEKGCNIRTQ SYKYRFAWGV VTCEPGPFCH VNLACDRKL VYCLKRNLRS YNPQYQYFPN ILCS.		
Solubility	Add 0.1M Acetate buffer pH4 to prepare a working stock solution of approximately 0.5 mg/mL and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10µg/ml. In higher concentrations the solubility of this antigen is limited. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.		
Formulation	Filtered (0.4um) and lyophilized in 0.5 mg/mL in 0.05M Acetate buffer pH4.		

Introduction

Phospholipase A2 (PLA2) catalyzes the hydrolysis of the sn-2 position of membrane glycerophospholipids to liberate arachidonic acid (AA), a precursor of eicosanoids including prostaglandins and leukotrienes. The same reaction also produces lysophospholipids, which represent another class of lipid mediators. The secretory PLA2 (sPLA2) family, in which 10 isozymes have been identified, consists of low molecular weight, Ca²⁺-requiring secretory enzymes that have been implicated in a number of biological processes, such as modification of eicosanoid generation, inflammation, and host defense. This enzyme has been proposed to hydrolyze phosphatidylcholine (PC) in

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

lipoproteins to liberate lyso-PC and free fatty acids in the arterial wall, thereby facilitating the accumulation of bioactive lipids and modified lipoproteins in atherosclerotic foci. In mice, sPLA2 expression significantly influences HDL particle size and composition and demonstrate that an induction of sPLA2 is required for the decrease in plasma HDL cholesterol in response to inflammatory stimuli. Instillation of bacteria into the bronchi was associated with surfactant degradation and a decrease in large:small ratio of surfactant aggregates in rats.

Stability

Store lyophilized protein at -20°C . Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C .

Background

Secreted Phospholipase A2-V Human Recombinant was produced with N-terminal His-Tag. PLA2G5 His-Tagged Fusion protein is 15.5 kDa containing 118 amino acid residues of the human secreted phospholipase A2-V and 16 additional amino acid residues - His-Tag (underlined).

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA