
Product Data Sheet

Product Name: B2M Human Recombinant
Cat. No.: GP22828
Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	Beta-2-microglobulin; B2M.		
Solubility	It is recommended to reconstitute the lyophilized B2M in sterile 18MΩ-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.		
Formulation	The protein was lyophilized from a concentrated solution (1mg/ml) containing PBS (pH 7.4) and 0.05% NaN3.		

Introduction

β2 microglobulin is an 11 kDa protein associated with the outer membrane of many cells including lymphocytes. It is the small subunit of the MHC class I molecule. Association with beta 2-microglobulin is generally required for the transport of class I heavy chains from the endoplasmic reticulum to the cell surface. β2 microglobulin associates with class I-like molecules such as CD1 and Qa as well as with the alpha chain of MHC class I molecules. Very limited amounts of MHC class I molecules can be found on the surface in the absence of β2 microglobulin. CD8 T cells cannot develop in the absence of MHC class I. Beta 2-microglobulin is present in small amounts in serum, csf, and urine of normal people, and to a much greater degree in the urine and plasma of patients with tubular proteinaemia, renal failure, or kidney transplants. Human Beta 2 microglobulin levels can rise either because its rate of synthesis has increased (e.g. in AIDS, malignant monoclonal plasma cell dyscrasia, solid tumors and autoimmune disease) or because of impaired renal filtration (e.g. due to renal insufficiency, graft rejection or nephrotoxicity induced by post-transplantation immunosuppressive therapy). Beta-2 microglobulin levels might also be elevated in multiple myeloma and lymphoma cases. Dialysis-related amyloidosis develops after a long-term hemodialysis, it can aggregate into amyloid fibers that deposit in joint spaces.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Stability

Lyophilized B2M although stable at room temperature for 3 weeks, should be stored desiccated below -18°C . Upon reconstitution B2M should be stored at 4°C between 2-7 days and for future use below -18°C . For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Background

B2 Microglobulin Human Recombinant produced in E.Coli is a non-glycosylated polypeptide chain having a molecular mass of 11.76 kDa. The B2M is purified by proprietary chromatographic techniques.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA