
Product Data Sheet

Product Name: CEA Human, His

Cat. No.: GP23109

Batch No.: 1

Product Data

Purity >98% Source Sf9, Baculovirus cells.

Physical Appearance solid Shipping Condition Shipped with Ice Packs.

Synonyms Carcinoembryonic Antigen Related Cell Adhesion Molecule 5; Carcinoembryonic Antigen-Related Cell Adhesion Molecule 5; Meconium Antigen 100; CEA; Carcinoembryonic Antigen; CD66e Antigen; CD66e; Carcinoembryonic antigen-related cell adhesion molecule 5; Carcinoembryonic antigen; CEA.

Amino Acid Sequence
 KLTIESTPFN VAEGKEVLLL VHNL PQHLFG YSWYKGERVD GNRQIIGYVI
 GTQQATPGPA YSGREIIPN ASLLIQNIIQ NDTGFYTLHV IKSDLVNEEA
 TGQFRVYPEL PKPSISSNNS KPVEDKDAVA FTCEPETQDA TYLWWVNNQS
 LPVSPRLQLS NGNRTLTLFN VTRNDTASYKCETQNPVSAR RSDSVILNVL
 YGPDAPTISP LNTSYRSGEN LNLSCHAASN PPAQYSWVFN GTFQQSTQEL
 FIPNITVNNS GSYTCQAHNS DTGLNRTT VT TITVYAEPK PFITSNNSNP
 VEDEDAVALT CEPEIQNTTY LWWVNNQSLP VSPRLQLSND NRTLTLTLLSVT
 RNDVGPYECGIQNKLSVDHS DPVILNVLYG PDDPTISPSY TYRPGVNLS
 LSCHAASNPP AQYSWLIDGN IQQHTQELFI SNITEKNSGL YTCQANNSAS
 GHSRTTVKTI TVSAELPKPS ISSNNSKPVE DKDAVAFTCE PEAQNTTYLW
 WVNGQSLPVS PRLQLSNGNR TLTLFNVTRN DARAYVCGIQNSVSANRSDP
 VTLDVLYGPD TPIISPPDSS YLSGANLNL CHSASNPSQ YSWRINGIPQ
 QHTQVLFI AK ITPNNGTYA CFVSNLATGR NNSIVKSITV SASGTSPGLS
 ALEHHHHHH.

Formulation CEA protein solution (0.5mg/ml) contains Phosphate Buffered Saline (pH 7.4) and 10% glycerol.

Introduction

Carcinoembryonic antigen (CEA) is a glycoprotein present in fetal digestive-tract tissues; it's involved in cell adhesion. The production of CEA stops before birth. CEA is called

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

tumor marker since its elevated levels are found in the serum from individuals with colorectal, gastric, pancreatic, lung and breast carcinomas and in heavy smokers. There are also benign conditions that elevate CEA levels such as smoking, infection, inflammatory bowel disease, pancreatitis, cirrhosis of the liver, and some benign tumors (in the equivalent organs which have cancers with elevated CEA). Typically, higher levels of CEA are found in men, smokers, and older individuals. The presence of CEA assists in screening, in evaluating recurrent or disseminated disease, and in determining the success of surgical removal of malignant tumors. CEA levels can be used as indicators of treatment success. The normal values range from 0.0 to 2.5 ng/ml of serum (from blood), in non-smokers, a greater amount than that may be suggestive of cancer. Levels above 20 ng/ml before treatment are associated with cancer which has already metastasized. Benign conditions do not usually cause a CEA increase over 10 ng/ml. The high levels of CEA should return to normal after successful therapy, however if during follow up there's an elevation in CEA levels it indicates a recurrence of tumor. Carcinoembryonic antigen family belongs to the immunoglobulin superfamily; it consists of 29 genes, 18 of which are normally expressed.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Background

CEA Human Recombinant produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 659 amino acids (35-685a.a.) and having a molecular mass of 72.3kDa (Molecular size on SDS-PAGE will appear at approximately 70-150kDa). CEA is expressed with an 8 amino acids His tag at C-Terminus and purified by proprietary chromatographic techniques.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA