
Product Data Sheet

Product Name: EIF3F Human
 Cat. No.: GP23381
 Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped with Ice Packs.
Synonyms	Eukaryotic Translation Initiation Factor 3; Subunit F; EIF3S5 ; Eukaryotic Translation Initiation Factor 3; Subunit 5 (Epsilon; 47kD); EC 3.4.19.12; Deubiquitinating Enzyme EIF3f; eIF3f; eIF-3-epsilon; EIF3 P47; Eukaryotic Translation Initiation Factor 3 Subunit 5; eIF3-p47; Eukaryotic Translation Initiation Factor 3; Subunit 5 Epsilon; 47kDa.		
Amino Acid Sequence	MGSSHHHHHH SGLVPRGSH MGSMATPAVP VSAPPATPTP VPAAAPASVP APTPAPAAAP VPAAAPASSS DPAAAAAATA APGQTPASAQ APAQTPAPAL PGPALPGPPF GGRVRLHPV ILASIVDSYE RRNEGAARVI GTLLGTVDKH SVEVTNCFVS PHNESEDEVA VDMEFAKNMY ELHKKVSPNE LILGWYATGH DITEHSVLII EYYSREAPNP IHLTVDTSLQ NGRMSIKAYV STLMGVPGRT MGVMFTPLTV KYAYYDTERI GVDLIMKTCF SPNRVIGLSS DLQQVGGASA RIQDALSTVL QYAEDVLSGK VSADNTVGRF LMSLVNQVPK IVPDDFETML NSNINDLLMV TYLANLTQSQ IALNEKLVNL.		
Formulation	EIF3F protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 10% glycerol, and 0.4M urea.		

Introduction

Eukaryotic Translation Initiation Factor 3F, also known as EIF3F belongs to the eIF-3 subunit F family. EIF3F is a component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is essential for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF 2: GTP: methionyl-tRNA_i and eIF-5 to form the 43S preinitiation complex (43S PIC). In addition, the eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also necessary for disassembly and recycling of posttermination ribosomal complexes and subsequently

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

prevents premature union of the 40S and 60S ribosomal subunits prior to initiation. Among the diseases associated with EIF3F are intrahepatic cholangiocarcinoma, and cholangiocarcinoma.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Background

EIF3F Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 380 amino acids (1-357 a.a) and having a molecular mass of 40kDa. EIF3F is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA