
Product Data Sheet

Product Name: SPA-Cys
 Cat. No.: GP24672
 Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped at Room temp.
Synonyms	Immunoglobulin G-binding protein A; IgG-binding protein A; Staphylococcal protein A; SPA.		
Amino Acid Sequence	<p>NAAQHDEAAQQ NAFYQVLNMP NLNADQRNGF IQSLKDDPSQ SANVLGEAQK LNSQAPKAD AQQNNFNKDQ QSAFYEILNM PNLNEAQRNG FIQSLKDDPS QSTNVLGEAK KLNESQAPKA DNNFNKEQQN AFYEILNMPN LNEEQRNGFI QSLKDDPSQS ANLLSEAKKL NESQAPKADN KFNKEQQNAF YEILHLPNLN EEQRNGFIQS LKDDPSQSAN LLAEAKKLND AQAPKADNKF NKEQQNAFYE ILHLPNLTEE QRNGFIQSLK DDPSVSKEIL AEAKKLNDAQ APKEEDC</p>		
Solubility	It is recommended to reconstitute the lyophilized SPA in sterile 18MΩ-cm H ₂ O not less than 0.1mg/ml, which can then be further diluted to other aqueous solutions.		
Formulation	SPA protein was lyophilized with no additives.		

Introduction

Protein A is a cell wall component produced by several strains of *Staphylococcus aureus*. The recombinant Protein A is genetically engineered protein and holds 5 IgG-binding regions of protein A. Recombinant Protein A functions basically the same as native Protein A and is perfect for purification of polyclonal or monoclonal IgG antibodies. Protein A binds to human IgG1, IgG2 and IgG4, mouse IgG2a, IgG2b and IgG3 and rat IgG2c. Protein A also binds to total IgG from rabbit, pig, dog, cat, and guinea pig.

Stability

Lyophilized SPA although stable at room temperature for 3 weeks, should be stored desiccated below -18°C . Upon reconstitution SPA should be stored at 4°C between 2-7 days and for future use below -18°C . For long term storage it is recommended to add a

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

carrier protein (0.1% HSA or BSA).

Background

SPA-Cys Recombinant produced in E.Coli is a single non-glycosylated polypeptide chain with a Cys on C-terminus. SPA-Cys is comprised of 5 IgG-binding domains E-D-A-B-C aligned in series containing 297 amino acids and having a molecular mass of 33.5kDa containing little or no carbohydrate. Cell wall binding region, cell membrane binding region and albumin binding region were removed to ensure the highest specific IgG binding.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA