
Product Data Sheet

Product Name: T.gondii p29
Cat. No.: GP25363
Batch No.: 1

Product Data

Purity	>98%	Source	Escherichia Coli.
Physical Appearance	solid	Shipping Condition	Shipped with Ice Packs.
Formulation	50mM Tris-HCl, 1.5M urea and 50% glycerol.		

Introduction

The life cycle of *Toxoplasma gondii* has two phases. The coccidia like takes place only in members of the Felidae family which makes these animals the parasite's primary host. The asexual part of the life cycle can take place in any warm-blooded animal, like other mammals(including felines) and birds. *T. gondii* constructing daughter scaffolds within the mother cell. In the intermediate hosts (including felines), the parasite invades cells, forming intracellular so-called parasitophorous vacuoles containing bradyzoites, the slowly replicating form of the parasit. Vacuoles form tissue cysts mainly within the muscles and brain. Since they are within cells, the host's immune system does not detect these cysts. Resistance to antibiotics varies, but the cysts are very difficult to eradicate entirely. Within these vacuoles *T. gondii* propagates by a series of binary fissions until the infected cell eventually bursts and tachyzoites are released. Tachyzoites are the motile, asexually reproducing form of the parasite. Unlike the bradyzoites, the free tachyzoites are usually efficiently cleared by the host's immune response, although some manage to infect cells and form bradyzoites, thus maintaining the infection.

Stability

Toxoplasma protein although stable at 4°C for 1 week, should be stored below -18°C .Please prevent freeze thaw cycles.

Background

The E.coli derived recombinant protein contains the p29 (GRA7) immunodominant regions, amino acids 24-100 and fused to a 26kDa GST tag.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA